

Normalization Properties of $\lambda\mu$ -Calculus Using Realizability Semantics

Péter Battyányi

University of Debrecen, Hungary

Karim Nour

Université Savoie Mont Blanc, France

The simply typed lambda calculus

The λ -terms:

$$\Lambda = \mathcal{V} \mid (\lambda \mathcal{V}. \Lambda) \mid (\Lambda) \Lambda,$$

where $\mathcal{V} = \{x, y, z, \dots\}$ are variables.

We adopt Krivine's notation: $(M)N$ stands for the application of N to M .

Moreover,

$$(M_1)M_2 \dots M_{n-1}M_n \rightleftharpoons (\dots (M_1)M_2) \dots M_{n-1})M_n.$$

The simply typed lambda calculus

The types are built from a set $\mathcal{V}_{\mathbb{T}}$ of atomic types with the connective $\rightarrow$:

$$\mathbb{T} := \mathcal{V}_{\mathbb{T}} \mid \mathbb{T} \rightarrow \mathbb{T}$$

Typing rules

A context Γ : a finite set of declarations of the form $x : A$.

Let $|\Gamma| = \{A \mid x : A \in \Gamma \text{ for some } x\}$. Then:

$$\overline{\Gamma, x : A \vdash x : A}$$

$$\frac{\Gamma, x : A \vdash M : B}{\Gamma \vdash \lambda x. M : A \rightarrow B}$$

$$\frac{\Gamma \vdash M : A \rightarrow B \quad \Gamma \vdash N : A}{\Gamma \vdash (M)N : B}$$

The λ -term M is called typable with type A , if $\Gamma \vdash M : A$ is derivable using the rules and axioms.

An example

Let $\Gamma = \{x : A \rightarrow (B \rightarrow C), y : A \rightarrow B, z : A\}$. Then,

$$\begin{array}{c}
 \frac{\overline{\Gamma \vdash x : A \rightarrow (B \rightarrow C)} \quad \overline{\Gamma \vdash z : A}}{\Gamma \vdash (x)z : B \rightarrow C} \quad \frac{\overline{\Gamma \vdash y : A \rightarrow B} \quad \overline{\Gamma \vdash z : A}}{\Gamma \vdash (y)z : B} \\
 \hline
 \Gamma \vdash ((x)z)(y)z : C \\
 \hline
 x : A \rightarrow (B \rightarrow C), y : A \rightarrow B \vdash \lambda z.((x)z)(y)z : A \rightarrow C \\
 \hline
 x : A \rightarrow (B \rightarrow C) \vdash \lambda yz.((x)z)(y)z : (A \rightarrow B) \rightarrow (A \rightarrow C) \\
 \hline
 \vdash \lambda xyz.((x)z)(y)z : (A \rightarrow (B \rightarrow C)) \rightarrow ((A \rightarrow B) \rightarrow (A \rightarrow C))
 \end{array}$$

Types as formulas

When we omit the indications of the terms, we obtain a correct derivation of the end formula.

$$\begin{array}{c}
 \frac{\overline{|\Gamma| \vdash A \rightarrow (B \rightarrow C)} \quad \overline{|\Gamma| \vdash A}}{\overline{|\Gamma| \vdash B \rightarrow C}} \quad \frac{\overline{|\Gamma| \vdash A \rightarrow B} \quad \overline{|\Gamma| \vdash A}}{\overline{|\Gamma| \vdash B}} \\
 \hline
 \overline{|\Gamma| \vdash C} \\
 \hline
 \overline{A \rightarrow (B \rightarrow C), A \rightarrow B \vdash A \rightarrow C} \\
 \hline
 \overline{A \rightarrow (B \rightarrow C) \vdash (A \rightarrow B) \rightarrow (A \rightarrow C)} \\
 \hline
 \vdash (A \rightarrow (B \rightarrow C)) \rightarrow ((A \rightarrow B) \rightarrow (A \rightarrow C))
 \end{array}$$

The Curry-Howard isomorphism

This phenomenon is referred to as the Curry-Howard isomorphism and was first observed by Curry in the 1930s and (re)discovered by Howard in 1968. It was widely believed that the Curry-Howard isomorphism is valid only for intuitionistic logic, until Griffin and Murthy discovered in the 1990s that certain terms expressing control can be given the type $\neg\neg A \rightarrow A$.

The classical case

From that point on, several calculi realizing the Curry-Howard isomorphism for classical logic have been discovered. For example,

- ▶ Rehof and Sorensen's λ_{Δ} ,
- ▶ Curien and Herbelin's $\bar{\lambda}\mu\tilde{\mu}$,
- ▶ Krivine's $\lambda\mathcal{C}$,
- ▶ Berardi and Barbanera's λ^{Sym} ,
- ▶ Parigot's $\lambda\mu$.

Definitions and notation

- ▶ Let $\mathcal{V}_\lambda = \{x, y, z, \dots\}$ denote a set of λ -variables and $\mathcal{V}_\mu = \{\alpha, \beta, \gamma, \dots\}$ denote a set of μ -variables, respectively. Formation rules for $\lambda\mu$ -terms:

$$\mathcal{T} \quad := \quad \mathcal{V}_\lambda \quad | \quad \lambda \mathcal{V}_\lambda. \mathcal{T} \quad | \quad (\mathcal{T}) \mathcal{T} \quad | \quad [\mathcal{V}_\mu] \mathcal{T} \quad | \quad \mu \mathcal{V}_\mu. \mathcal{T}$$

- ▶ In a $\lambda\mu$ -term, the λ and μ operators bind the variables in their scope. We consider terms modulo correct renaming of bound variables.

Definitions and notation

The types are built from a set $\mathcal{V}_{\mathbb{T}}$ of atomic types and the constant $\perp$ with the connectives $\neg$ and $\rightarrow$.

$$\mathbb{T} \quad := \quad \mathcal{V}_{\mathbb{T}} \cup \{\perp\} \quad | \quad \neg \mathbb{T} \quad | \quad \mathbb{T} \rightarrow \mathbb{T}$$

A context: a pair (Γ, Θ) .

- ▶ Γ : finite set of declarations of the form $x : A$
- ▶ Θ : finite set of declarations of the form $\alpha : \neg A$

If $x : A \rightarrow \perp$, we may also write $\neg A$ for $A \rightarrow \perp$.

Definitions and notation

The typing rules are:

$$\overline{\Gamma, x : A; \Theta \vdash x : A} \text{ ax}$$

$$\frac{\Gamma, x : A; \Theta \vdash M : B}{\Gamma; \Theta \vdash \lambda x. M : A \rightarrow B} \rightarrow_i$$

$$\frac{\Gamma; \Theta \vdash M : A \rightarrow B \quad \Gamma; \Theta \vdash N : A}{\Gamma; \Theta \vdash (M)N : B} \rightarrow_e$$

$$\frac{\Gamma; \alpha : \neg A, \Theta \vdash M : A}{\Gamma; \alpha : \neg A, \Theta \vdash [\alpha]M : \perp} \perp_i$$

$$\frac{\Gamma; \alpha : \neg A, \Theta \vdash M : \perp}{\Gamma; \Theta \vdash \mu \alpha. M : A} \perp_e$$

Definitions and notation

Below, let $\Gamma = \{x : \neg B \rightarrow \neg A, y : A, z : B\}$, $\Theta = \{\alpha : \neg B\}$ and $\Gamma' = \Gamma \setminus \{z : B\}$. Then

$$\begin{array}{c}
 \frac{\Gamma; \Theta \vdash z : B}{\Gamma; \Theta \vdash [\alpha]z : \perp} \\
 \frac{\Gamma'; \Theta \vdash \lambda z. [\alpha]z : \neg B}{\Gamma'; \Theta \vdash (x)\lambda z. [\alpha]z : \neg A} \quad \Gamma'; \Theta \vdash y : A \\
 \frac{\Gamma'; \Theta \vdash ((x)\lambda z. [\alpha]z)y : \perp}{\Gamma' \vdash \mu\alpha. ((x)\lambda z. [\alpha]z)y : B} \\
 \frac{x : \neg B \rightarrow \neg A \vdash \lambda y. \mu\alpha. ((x)\lambda z. [\alpha]z)y : A \rightarrow B}{\vdash \lambda xy. \mu\alpha. ((x)\lambda z. [\alpha]z)y : (\neg B \rightarrow \neg A) \rightarrow (A \rightarrow B)}
 \end{array}$$

Definitions and notation

If we omit the terms annotating the type formulas, we obtain a full fledged proof of $(\neg B \rightarrow \neg A) \rightarrow (A \rightarrow B)$.

$$\begin{array}{c}
 \frac{\frac{\frac{| \Gamma | \vdash B^{(1)} \quad | \Gamma | \vdash \neg B^{(2)}}{| \Gamma | \vdash \perp}}{| \Gamma' | \vdash \neg B} \quad (1)}{| \Gamma' | \vdash \neg B \rightarrow \neg A^{(4)}} \quad \frac{}{| \Gamma' | \vdash \neg A} \quad \frac{}{| \Gamma' | \vdash A^{(3)}} \\
 \hline
 \frac{}{| \Gamma' | \vdash \perp} \quad (2) \\
 \frac{}{\neg B \rightarrow \neg A, A \vdash B} \quad (3) \\
 \frac{}{\neg B \rightarrow \neg A \vdash A \rightarrow B} \quad (4) \\
 \hline
 \vdash (\neg B \rightarrow \neg A) \rightarrow (A \rightarrow B)
 \end{array}$$

Reduction rules

We use several types of substitution in the $\lambda\mu$ -calculus.

- ▶ β -substitution: $M[x := N]$ is obtained from M by exchanging the free occurrences of x in M with N .
- ▶ μ -substitution: $M[\alpha :=_r N]$ is obtained from M when we replace all subterm occurrences $[\alpha]U$ of M by $[\alpha](U)N$ recursively.
- ▶ μ' -substitution: $M[\alpha :=_l N]$ is obtained from M when we replace all subterm occurrences $[\alpha]U$ of M by $[\alpha](N)U$ in a recursive manner.
- ▶ ρ -substitution: $M[\alpha := \beta]$ is obtained from M by replacing the free occurrences of α in M by β

The notation M_α stands for the erasure of every occurrence of $[\alpha]$ from M .

Reduction rules

With this in hand, we are able to define the reduction rules.

- ▶ β -reduction: $(\lambda x^A.M^B)N^A : B \rightarrow_\beta M^B[x := N] : B$
- ▶ μ -reduction: $(\mu\alpha^{\neg(A \rightarrow B)}.M^\perp)N^A : B \rightarrow \mu\alpha^{\neg B}.M^\perp[\alpha :=_r N] : B$
- ▶ μ' -reduction: $(N)^{A \rightarrow B} \mu\alpha^{\neg A}.M^\perp : B \rightarrow \mu\alpha^{\neg B}.M^\perp[\alpha :=_l N] : B$
- ▶ ρ -reduction: $[\alpha^{\neg A}] \mu\beta^{\neg A}.M^\perp : \perp \rightarrow_\rho M^\perp[\beta := \alpha] : \perp$
- ▶ θ -reduction: $\mu\alpha^{\neg A}.[\alpha^{\neg A}]M^A : A \rightarrow_\theta M^A : A$ provided $\alpha \notin fv(M)$
- ▶ ε -reduction: $\mu\alpha^{\neg A}.\mu\beta^{\neg \perp}.M^\perp : A \rightarrow_\varepsilon \mu\alpha^{\neg A}.M^\perp_\beta : A.$

Reduction rules

- β -reduction: eliminates subsequent occurrences of $\rightarrow_i$ and $\rightarrow_e$.

$$\frac{\frac{\Gamma, x : A; \Theta \vdash M : B}{\Gamma; \Theta \vdash \lambda x. M : A \rightarrow B} \rightarrow_i \quad \Gamma \vdash N : A}{\Gamma; \Theta \vdash (\lambda x. M)N : B} \rightarrow_e \rightsquigarrow \Gamma; \Theta \vdash M[x := N] : B$$

where $\Gamma \vdash M[x := N] : B$ is obtained from $\Gamma \vdash M : B$ by exchanging all hypotheses $x : A$ with the derivation $\Gamma \vdash N : A$.

Reduction rules

- ▶ μ -reduction: μ propagates an argument toward occurrences marked by α .
- ▶ Given the declaration $\alpha : A \rightarrow B$, we have $M : \perp$. We consider the declaration $\alpha : B$ and replace in M each subterm of the form $[\alpha]P$ by $[\alpha](P)N$ to obtain a term of type B .
- ▶ At the proof level, μ -reduction corresponds to the simplification of a reasoning by contradiction, when an application of a $\perp$ -elimination is replaced by $\perp$ -eliminations of lower type complexity.

Reduction rules

$$\frac{\frac{\Gamma; \alpha : \neg(A \rightarrow B), \Theta \vdash M : \perp}{\Gamma; \Theta \vdash \mu\alpha.M : A \rightarrow B} \perp_e \quad \Gamma; \Theta \vdash N : A}{\Gamma; \Theta \vdash (\mu\alpha.M)N : B} \rightarrow_e \rightsquigarrow \Gamma; \Theta \vdash \mu\alpha.M[\alpha :=_r N] : B$$

Locally,

$$[\alpha^{\neg(A \rightarrow B)}]U^{A \rightarrow B} \rightsquigarrow [\alpha^{\neg B}](U^{A \rightarrow B})N^A$$

Reduction rules

- ▶ μ' -reduction: μ' propagates the right-side functional part toward occurrences marked by an application with the μ -variable
- ▶ It is the symmetric counterpart of rule μ . Given the declaration $\alpha : \neg A$, we have $\mu\alpha.M : A$ by an application of $\perp_e$. If we also have $N : A \rightarrow B$, then $(N)\mu\alpha.M : B$ and $(N)\mu\alpha.M : B \rightarrow \mu\alpha.M[\alpha :=_I N]$. It suffices to start with the declaration $\alpha : B$ and to replace in M each subterm of the form $[\alpha]P$ by $[\alpha](N)P$.
- ▶ Here, we observe that there is no direct relation between the two types associated with α before and after the reduction.

Reduction rules

$$\frac{\Gamma; \Theta \vdash N : A \rightarrow B \quad \frac{\Gamma; \alpha : \neg A, \Theta \vdash M : \perp}{\Gamma; \Theta \vdash \mu\alpha.M : A} \perp_e}{\Gamma; \Theta \vdash (N)\mu\alpha.M : B} \rightarrow_e \rightsquigarrow \Gamma; \Theta \vdash \mu\alpha.M[\alpha :=_I N] : B$$

Locally,

$$[\alpha^{\neg A}]U^A \rightsquigarrow [\alpha^{\neg B}](N^{A \rightarrow B})U^A$$

Simplification rules

- ▶ ρ -reduction: If we obtained a derivation of $[\beta]\mu\alpha.M : \perp$, then, necessarily, $\alpha, \beta : \neg A$ for some A . Hence, ρ -reduction can be interpreted as a simplification of proofs: it is enough to replace every occurrence of the assumption $\alpha : \neg A$ by the same type of assumption $\beta : \neg A$ throughout the derivation.
- ▶ θ -reduction: Given the declaration $\alpha : \neg A$, we have $M : A$, hence $\mu\alpha.[\alpha]M : A$. This term reduces simply to M , provided that α does not appear free in M .
- ▶ ε -reduction: Given the declaration $\alpha : \neg A$ and the relation $\mu\alpha.\mu\beta.M : A$, we necessarily have $\beta : \neg\perp$. We eliminate $\mu\beta$ as well as all occurrences of $[\beta]$ in M to obtain a term of type A . Hence, ε -rule removes a proof of $\perp$ obtained from $\perp$ by contradiction.

Why the μ' -rule?

Proposition

Let t be a (closed) term of the second order propositional λ -calculus. Then

$$t : \mathcal{N} = X \rightarrow (X \rightarrow X) \rightarrow X \quad \text{iff} \quad t = \lambda x \lambda f. (f)^n x$$

for some $n \in \mathbb{N}$.

On the other hand,

$$\lambda x \lambda f. \mu \alpha [\alpha](f) \mu \beta [\alpha](f) \mu \gamma [\beta](f)(f) \mu \beta [\gamma](f) \mu \gamma [\beta] x$$

is a $\lambda\mu$ -term of type $\mathcal{N}$ with value 3.

Why the μ' -rule?

In Parigot's original calculus, a proof that n is a natural number may normalize to a term different from the Church numeral $\underline{n}$.

- ▶ Such terms are called *classical natural numbers*.
- ▶ Their effective numerical value may be difficult to recover.

Possible solutions

- ▶ An algorithm for extracting the value of a classical natural number (Parigot).
- ▶ A Krivine-type storage operator to simulate call-by-value in a call-by-name language (Nour).
- ▶ The introduction of the symmetric counterpart of μ as a new rule, which is called the μ' -rule (Parigot).

Reduction rules

Theorem (B.; Nour 2022)

Let x, f be two different λ -variables. If $\vdash M : \mathcal{N}$ and $((M)x)f \in \mathcal{WN}_{\beta\mu\mu'\rho\theta\varepsilon}$, then $\exists n \in \mathbb{N}: ((M)x)f \rightarrow_{\beta\mu\mu'\rho\theta\varepsilon} (f)^n x$.

In particular, to extract the value of a $\lambda\mu$ -term of type $\mathcal{N}$, we have make use of all the reductions $\beta\mu\mu'\rho\theta\varepsilon$.

Reduction rules

Let $M = \lambda x. \mu \alpha. [\alpha] \lambda g. (g) \mu \beta. \mu \gamma. [\alpha] \lambda f. (f) x$. Let $x : X$, $f : X \rightarrow X$, $g : X \rightarrow X$, $\alpha : \neg((X \rightarrow X) \rightarrow X)$, $\beta : \neg X$ and $\gamma : \neg \perp$, then $\vdash M : \mathcal{N}$. Furthermore,

$$\begin{aligned}
 ((M)x)f &= \underbrace{((\lambda x. \mu \alpha. [\alpha] \lambda g. (g) \mu \beta. \mu \gamma. [\alpha] \lambda f. (f) x)x)f}_{\rightarrow_{\beta}} \\
 &\rightarrow_{\beta} \underbrace{(\mu \alpha. [\alpha] \lambda g. (g) \mu \beta. \mu \gamma. [\alpha] \lambda f. (f) x)f}_{\rightarrow_{\mu}} \\
 &\rightarrow_{\mu} \mu \alpha. [\alpha] (\lambda g. (g) \mu \beta. \mu \gamma. [\alpha] \underbrace{(\lambda f. (f) x)f}_{\rightarrow_{\beta}}) f \\
 &\rightarrow_{\beta} \mu \alpha. [\alpha] \underbrace{(\lambda g. (g) \mu \beta. \mu \gamma. [\alpha] (f) x)f}_{\rightarrow_{\beta}} \\
 &\rightarrow_{\beta} \mu \alpha. [\alpha] (f) \underbrace{\mu \beta. \mu \gamma. [\alpha] (f) x}_{\rightarrow_{\varepsilon}} \\
 &\rightarrow_{\varepsilon} \mu \alpha. [\alpha] (f) \mu \beta. [\alpha] (f) x \\
 &\rightarrow_{\mu'} \mu \alpha. [\alpha] \underbrace{\mu \beta. [\alpha] (f) x}_{\rightarrow_{\rho}} \\
 &\rightarrow_{\rho} \mu \alpha. [\alpha] (f) x \\
 &\rightarrow_{\theta} (f) x.
 \end{aligned}$$

Why de Groote's syntax?

The typing rule representing reductio ad absurdum by Parigot:

$$\frac{\Gamma; \alpha : \neg A, \Theta \vdash M : A}{\Gamma; \Theta \vdash \mu\alpha.[\alpha]M : \perp} \perp_i$$

De Groote split it into two rules:

$$\frac{\Gamma; \alpha : \neg A, \Theta \vdash M : A}{\Gamma; \alpha : \neg A, \Theta \vdash [\alpha]M : \perp} \perp_i$$

$$\frac{\Gamma; \alpha : \neg A, \Theta \vdash M : \perp}{\Gamma; \Theta \vdash \mu\alpha.M : A} \perp_e$$

Why de Groote's syntax?

In Parigot's original calculus (when

$\Gamma = \{x : A, y : \neg\neg A\}, \Theta = \{\alpha : \neg A, \beta : \neg\perp, \gamma : \neg\perp\}$):

$$\begin{array}{c}
 \frac{\Gamma; \Theta \vdash x : A}{\Gamma; \Theta \vdash [\alpha]x : \perp} \\
 \frac{\Gamma; \Theta \setminus \{\beta : \neg\perp\} \vdash \mu\beta.[\alpha]x : \perp}{\Gamma \setminus \{x : A\}; \Theta \setminus \{\beta : \neg\perp\} \vdash \lambda x. \mu\beta.[\alpha]x : \neg A} \\
 \frac{\Gamma \setminus \{x : A\}; \Theta \setminus \{\beta : \neg\perp\} \vdash \lambda x. \mu\beta.[\alpha]x : \neg A}{\Gamma \setminus \{x : A\}; \Theta \setminus \{\beta : \neg\perp\} \vdash (y)\lambda x. \mu\beta.[\alpha]x : \perp} \\
 \frac{\Gamma \setminus \{x : A\}; \Theta \setminus \{\beta : \neg\perp\} \vdash (y)\lambda x. \mu\beta.[\alpha]x : \perp}{\Gamma \setminus \{x : A\}; \Theta \setminus \{\beta : \neg\perp\} \vdash [\gamma](y)\lambda x. \mu\beta.[\alpha]x : \perp} \\
 \frac{y : \neg\neg A; \gamma : \neg\perp \vdash \mu\alpha.[\gamma](y)\lambda x. \mu\beta.[\alpha]x : A}{\gamma : \neg\perp \vdash \lambda y. \mu\alpha.[\gamma](y)\lambda x. \mu\beta.[\alpha]x : \neg\neg A \rightarrow A}
 \end{array}$$

The hypothesis $\gamma : \neg\perp$ never disappears from the set of assumptions.

Why de Groote's syntax?

Possible solutions

- ▶ de Groote's modification of the syntax.
- ▶ λ -top calculus due to Herbelin and Ariola.

In what follows, we choose de Groote's version of the $\lambda\mu$ -calculus.

Why de Groote's syntax?

The previous proof in de Groote's terms (assuming $\Gamma = \{x : A, y : \neg\neg A\}$, $\Theta = \{\alpha : \neg A\}$):

$$\begin{array}{c}
 \Gamma; \Theta \vdash x : A \\
 \hline
 \Gamma; \Theta \vdash [\alpha]x : \perp \\
 \hline
 \Gamma \setminus \{x : A\}; \Theta \vdash \lambda x. [\alpha]x : \neg A \\
 \hline
 \Gamma \setminus \{x : A\}; \Theta \vdash (y)\lambda x. [\alpha]x : \perp \\
 \hline
 y : \neg\neg A \vdash \mu\alpha. (y)\lambda x. [\alpha]x : A \\
 \hline
 \vdash \lambda y. \mu\alpha. (y)\lambda x. [\alpha]x : \neg\neg A \rightarrow A
 \end{array}$$

Two calculi, two normalization problems

We study two reduction systems:

Without μ'

$\beta\mu\rho\varepsilon\theta$

With μ'

$\beta\mu\mu'\rho\varepsilon\theta$

Strong vs. weak normalization

Definition

- 1 $M \in \mathcal{SN}_{\mathcal{R}}$ if every $\mathcal{R}$ -reduction sequence starting from M is finite.
- 2 $M \in \mathcal{WN}_{\mathcal{R}}$ if *some* $\mathcal{R}$ -reduction sequence starting from M reaches a normal form.

Theorem

- 1 (Parigot, 1993) Let $\mathcal{R} = \beta\mu\rho\varepsilon\theta$ in the typed $\lambda\mu$ -calculus. Then $\mathcal{R}$ is strongly normalizing.
- 2 (B.; Nour, 2022) Let $\mathcal{R} = \beta\mu\mu'\rho\varepsilon\theta$ in the simply typed $\lambda\mu$ -calculus. Then $\mathcal{R}$ is weakly normalizing.

We remark that $\mu\mu'$ together with ρ is not strongly normalizing and together with ε is not even weakly normalizing (B., 2007).

Why a new proof method?

Earlier normalization proofs were mainly syntactic and arithmetical.

They typically required

- ▶ inductions on several measures;
- ▶ separate treatment of different reduction rules;
- ▶ technical commutation lemmas;
- ▶ substantial reconstruction when new reductions were added.

Goal

Develop a single semantic framework in which the desired normalization property follows by choosing an appropriate parameter for the interpretation.

The realizability strategy

The basic idea is to interpret each type A by a set of terms $\mathcal{I}(A)$, where $\mathcal{I}(A) \subseteq \perp\!\!\!\perp$.

The main goal is:

$$\Gamma; \Theta \vdash M : A \quad \Longrightarrow \quad M \in \mathcal{I}(A).$$

This would imply

$$M \text{ is typable} \quad \Longrightarrow \quad M \in \perp\!\!\!\perp.$$

Thus, by assigning different computational meanings to $\perp\!\!\!\perp$, we obtain different properties of the calculus. Hence, the semantics is parametrized by $\perp\!\!\!\perp$.

The realizability strategy¹

The case of the simply typed lambda calculus with $\perp = SN_\beta$ (Tait, 1967; Girard, 1971):

Definition

Let Λ denote the simply typed λ -calculus terms.

1 If $\mathcal{A}, \mathcal{B} \subseteq \Lambda$, then

$$\mathcal{A} \rightarrow \mathcal{B} = \{F \in \Lambda \mid (\forall a \in \mathcal{A})(Fa \in \mathcal{B})\}.$$

2 $\llbracket A \rrbracket = \perp (= SN_\beta)$ if $A \in \mathcal{V}_\mathbb{T}$.

3 $\llbracket X \rrbracket = \llbracket U \rrbracket \rightarrow \llbracket V \rrbracket$, where $X = U \rightarrow V$.

¹Based on Sørensen; Urzyczyn: Lectures on the Curry-Howard isomorphism

The realizability strategy

Definition

1 A set $\mathcal{S} \subseteq \mathbb{L}$ is *saturated* if

① For all $n \geq 0$ and $M_1, \dots, M_n \in \mathbb{L}$:

$$(x) M_1 \cdots M_n \in \mathcal{S}$$

② For all $n \geq 1$ and $M_1, \dots, M_n \in \mathbb{L}$

$$(M_0[x := M_1]) M_2 \cdots M_n \in \mathcal{S} \Rightarrow (\lambda x. M_0) M_1 M_2 \cdots M_n \in \mathcal{S}$$

2

$$\mathbb{S} = \{X \subseteq \Lambda \mid X \text{ is saturated}\}.$$

The realizability strategy

Lemma

- 1 $\perp (= \mathcal{SN}_\beta) \in \mathbb{S}$;
- 2 $\mathcal{A}, \mathcal{B} \in \mathbb{S} \Rightarrow \mathcal{A} \rightarrow \mathcal{B} \in \mathbb{S}$;
- 3 $X \in \Pi \Rightarrow \llbracket X \rrbracket \in \mathbb{S}$. In particular, if $x : X$, then $x \in \llbracket X \rrbracket$.

The realizability strategy

Definition

- 1 A valuation is a map $\rho : \mathcal{V} \rightarrow \Lambda$, where $\mathcal{V}$ is the set of term variables. The valuation $\rho(x := N)$ is defined by

$$\rho(x := N)(y) = \begin{cases} N & \text{if } x = y, \\ \rho(y) & \text{otherwise.} \end{cases}$$

- 2 Let ρ be a valuation. Then

$$\llbracket M \rrbracket_\rho = M[x_1 := \rho(x_1), \dots, x_n := \rho(x_n)],$$

where $\text{FV}(M) = \{x_1, \dots, x_n\}$.

The realizability strategy

Definition

3 Let ρ be a valuation. Then

$$\rho \models M : A \quad \text{iff} \quad \llbracket M \rrbracket_\rho \in \llbracket A \rrbracket.$$

Also, $\rho \models \Gamma$ iff $\rho(x) \in \llbracket A \rrbracket$ for all $x : A \in \Gamma$.

4

$$\Gamma \models M : A \quad \text{iff} \quad \forall \rho : \rho \models \Gamma \Rightarrow \rho \models M : A.$$

The realizability strategy

Lemma (Adequacy)

$$\Gamma \vdash M : A \implies \Gamma \models M : A.$$

Theorem

$$\Gamma \vdash M : A \implies M \in \text{SN}_\beta.$$

Proof.

If $\Gamma \vdash M : A$, then $\Gamma \models M : A$. For each $x : X \in \Gamma$, let $\rho(x) = x$. Then $x \in \llbracket X \rrbracket$ holds since $\llbracket X \rrbracket \in \mathbb{S}$. Then $\rho \models \Gamma$, and we have

$$M = \llbracket M \rrbracket_\rho \in \llbracket A \rrbracket \subseteq \text{SN}_\beta.$$

The case of the $\lambda\mu$ -calculus

The main difference lies in the following typing rules:

$$\begin{aligned}\alpha : \neg A, M : \perp &\implies [\alpha]M : \perp, \\ M : \perp, \alpha : \neg A &\implies \mu\alpha.M : A..\end{aligned}$$

If we interpret $\llbracket \perp \rrbracket = \perp$, the interpretation must allow us to pass from

$$M \in \perp \quad \text{to} \quad \mu\alpha.M \in \llbracket A \rrbracket.$$

Crucial idea (Parigot, 1997)

For a set $\mathcal{X}$ of finite sequences of terms in $\mathbb{L}$, define

$$\mathcal{X} \rightsquigarrow \mathbb{L} = \{M \mid (M)\bar{P} \in \mathbb{L} \text{ for any initial segment } \bar{P} \text{ of an arbitrary element of } \mathcal{X}\}.$$

Intuitively:

- ▶ $\mathcal{X}$ describes a collection of finite sequences of arguments,
- ▶ $M \in \mathcal{X} \rightsquigarrow \mathbb{L}$ means that M sends all such sequences into $\mathbb{L}$.

This is the operation used to construct the reducibility candidates.

Saturation

A saturated set $\mathcal{S} \subseteq \Lambda$ is stable under term construction and expansion with β - and μ .
In our case:

$$\begin{aligned}
 M \in \mathcal{S}, x \in \mathcal{V}_\lambda &\implies \lambda x.M \in \mathcal{S}, \\
 M \in \mathcal{S}, \alpha \in \mathcal{V}_\mu &\implies \mu \alpha.M \in \mathcal{S}, \\
 M \in \mathcal{S}, \alpha \in \mathcal{V}_\mu &\implies [\alpha]M \in \mathcal{S}, \\
 x \in \mathcal{V}_\lambda, \bar{N} \in \mathbb{L}^{<\omega} &\implies (x)\bar{N} \in \mathcal{S}, \\
 M \in \mathcal{T}, \bar{P} \in \mathcal{T}^{<\omega}, N \in \mathcal{S}, (M[x := N])\bar{P} \in \mathcal{S} &\implies (\lambda x.M)N\bar{P} \in \mathcal{S}, \\
 M \in \mathcal{T}, \bar{N} \in \mathcal{S}^{<\omega}, \mu \alpha.M[\alpha :=_r \bar{N}] \in \mathcal{S} &\implies (\mu \alpha.M)\bar{N} \in \mathcal{S}.
 \end{aligned}$$

Semantic consequence

Definition (The $\rightsquigarrow$ operator)

Let $\perp$ be a saturated set and consider $\mathcal{K}, \mathcal{L} \subseteq \perp$, and $\mathcal{X} \subseteq \perp^{<\omega}$. Then:

$$\mathcal{K} \rightsquigarrow_{\perp} \mathcal{L} = \{M \in \perp \mid \forall N \in \mathcal{K}, (M)N \in \mathcal{L}\},$$

$$\mathcal{X} \rightsquigarrow_{\perp} \perp = \{M \in \perp \mid \forall \bar{N} \in \mathcal{X}, \text{ if } \bar{P} \sqsubseteq \bar{N}, \text{ then } (M)\bar{P} \in \perp\},$$

where $\bar{P} \sqsubseteq \bar{N}$ means that $\bar{P}$ is an initial segment of $\bar{N}$.

Definition ($\perp$ -saturated sets)

Let $\perp$ be saturated. We call $\mathcal{S}$ a $\perp$ -saturated set if the following property holds:

$$\exists \mathcal{X}_{\mathcal{S}} \subseteq \perp^{<\omega}, \mathcal{S} = \mathcal{X}_{\mathcal{S}} \rightsquigarrow_{\perp} \perp.$$

Weakest precondition

For a semantic set $\mathcal{S}$, define its weakest precondition

$$\mathcal{S}^{\perp} = \bigcup \{ \mathcal{X} \subseteq \perp^{<\omega} \mid \mathcal{S} = \mathcal{X} \rightsquigarrow \perp \}.$$

The crucial property is

$$\mathcal{S} = \mathcal{S}^{\perp} \rightsquigarrow \perp.$$

Interpretation

- ▶ We define a $\perp$ -model $\mathcal{M}$ in a way such that, for every $S \in \mathcal{M}$, S is $\perp$ -saturated.
- ▶ If $\mathcal{M}$ is a $\perp$ -model, then an $\perp$ -interpretation $\mathcal{I}$ is a function such that
 - $\mathcal{I}(\perp) = \perp$,
 - $\mathcal{I}(A \rightarrow B) = \mathcal{I}(A) \rightsquigarrow \mathcal{I}(B)$.

These conditions are tailored so that the interpretation of every type remains inside $\perp$. That is, $\mathcal{I}(A) \subseteq \perp$ for any type A .

The adequacy theorem

Let $M \in \mathcal{T}$ and σ be the simultaneous substitution $[(x_i := M_i)_{1 \leq i \leq n}; (\alpha_j :=_r \bar{N}_j)_{1 \leq j \leq m}]$. Then $M\sigma$ is obtained from the $\lambda\mu$ -term M by replacing each x_i by M_i and replacing inductively each subterm of the form $[\alpha_j]U$ in M by $[\alpha_j](U)\bar{N}_j$.

Theorem (General correctness theorem)

Let $\mathcal{M}$ be a $\perp$ -model, $\mathfrak{I}$ an $\mathcal{M}$ -interpretation,
 $\Gamma = \{x_i : A_i\}_{1 \leq i \leq n}$, $\Theta = \{\alpha_j : \neg B_j\}_{1 \leq j \leq m}$,
 $M_i \in \mathfrak{I}(A_i)$ for all $1 \leq i \leq n$, $\bar{P}_j \sqsubseteq \bar{N}_j \in \mathfrak{I}(B_j)^\perp$ for all $1 \leq j \leq m$ and
 $\sigma = [(x_i := M_i)_{1 \leq i \leq n}; (\alpha_j :=_r \bar{P}_j)_{1 \leq j \leq m}]$.
 If $\Gamma; \Theta \vdash M : A$, then $M\sigma \in \mathfrak{I}(A)$.

The adequacy theorem

Proof.

By induction on the derivation $\Gamma; \Theta \vdash M : A$.

We consider only the case $\perp_e$. In this case, $\Gamma; \Theta \vdash M = \mu\alpha.N$, hence, $\Gamma; \Theta', \alpha : \neg A \vdash N : \perp$, where $\Theta = \Theta' + (\alpha : A)$.

- ▶ Since $\emptyset \in \mathcal{I}(A)^\perp$ and $N\sigma = N\sigma'$ where $\sigma' = \sigma + [\alpha :=_r \emptyset]$, we obtain, by IH, $N\sigma \in \mathcal{I}(\perp) = \perp$, hence, $M\sigma = \mu\alpha.N\sigma \in \perp$.
- ▶ Let $\bar{P} \sqsubseteq \bar{N} \in \mathcal{I}(A)^\perp$. We have $(M\sigma)\bar{P} = (\mu\alpha.N\sigma)\bar{P} \rightarrow_\mu \mu\alpha.N\sigma''$ where $\sigma'' = \sigma + [\alpha :=_r \bar{P}]$. By IH, $N\sigma'' \in \perp = \mathcal{I}(\perp)$ and $\mu\alpha.N\sigma'' = \mu\alpha.N\sigma[\alpha :=_r \bar{P}] \in \perp$, hence $(M\sigma)\bar{P} \in \perp$. Therefore $M\sigma = \mu\alpha.N\sigma \in \mathcal{I}(A)$.

Correctness

Corollary (Correctness lemma)

If $\Gamma; \Theta \vdash M : A$ and $\perp$ is a saturated set, then $M \in \perp$.

Proof.

Assume $\Gamma = \{x_i : A_i\}_{1 \leq i \leq n}$ and $\Theta = \{\alpha_j : \neg B_j\}_{1 \leq j \leq m}$. Let $\mathcal{M}$ be the $\perp$ -model containing only the set $\perp$ and $\mathfrak{I}$ the $\mathcal{M}$ -interpretation defined by $\mathfrak{I}(X) = \perp$ for all $X \in \mathcal{V}_{\mathbb{T}}$. Since $x_i \in \mathfrak{I}(A_i)$ for all $1 \leq i \leq n$ and $\emptyset \in \mathfrak{I}(B_j)^\perp$ for all $1 \leq j \leq m$, then, by the general correctness lemma,

$M = M[(x_i := x_i)_{1 \leq i \leq n}; (\alpha_j :=_r \emptyset)_{1 \leq j \leq m}] \in \mathfrak{I}(A) \subseteq \perp$. □

The Parameter $\perp$

The semantics is parameterized by the set of terms $\perp$. Different choices of $\perp$ will yield different consequences.

To prove a normalization property for all typable terms, it suffices to prove that the corresponding set $\perp$ is saturated.

Case I: Strong normalization

Take

$$\perp\!\!\!\perp = \mathcal{T}_t \cap \mathcal{SN}_{\beta\mu\rho\varepsilon\theta}.$$

The main technical work is to verify the six saturation conditions.

Result

The set $\perp\!\!\!\perp$ is saturated.

Hence, by the correctness lemma,

$$M \in \mathcal{T}_t \implies M \in \mathcal{SN}_{\beta\mu\rho\varepsilon\theta}.$$

Why μ' changes the picture

Reductions μ , μ' and ρ together destroy strong normalization.

Let

$$U = \mu\alpha.[\alpha][\alpha]x, \quad M = (\mu\beta.U)U.$$

Taking $x : \perp$, $\alpha : \neg\perp$, $\beta : \neg(\perp \rightarrow \perp)$, M is typable, but there is a reduction sequence

$$\begin{aligned} M &\longrightarrow_{\mu'} \mu\alpha.[\alpha](\mu\beta.U)[\alpha](\mu\beta.U)x \longrightarrow_{\mu} \mu\alpha.[\alpha](\mu\beta.U)[\alpha]\mu\beta.U \\ &\longrightarrow_{\rho} \mu\alpha.[\alpha]M(\longrightarrow_{\theta} M). \end{aligned}$$

Therefore

$$M \notin \mathcal{T}_t \cap \mathcal{SN}_{\beta\mu\mu'\rho\in\theta}.$$

Case II: Weak normalization

The case of $\beta\mu\mu'\rho\epsilon\theta$ -reduction must concern *weak*, rather than strong, normalization. Now take

$$\mathbb{L}' = \mathcal{T}_t \cap \mathcal{WN}_{\beta\mu\mu'\rho\epsilon\theta}.$$

Result

$\mathbb{L}'$ is saturated.

Hence,

$$M \in \mathcal{T}_t \implies M \in \mathcal{WN}_{\beta\mu\mu'\rho\epsilon\theta}.$$

In both cases, the assumption of typability is essential for the proofs to work.

Summary

Advantages

- ▶ All reduction rules are treated simultaneously. Logical (β, μ, μ') and simplification rules $(\rho, \varepsilon, \theta)$ need not be separated.
- ▶ The parameter $\perp$ makes the proof “reusable”.

Disadvantage

Unlike the “combinatorial” proofs, realizability proofs do not allow us to extract normalization algorithms from the proofs.